

1885

1884-1886

William Osler first used the term “**mycotic aneurysm**” in **1885** to describe a mushroom-shaped aneurysm in a patient with subacute bacterial endocarditis ¹⁾.

Meralgia paresthetica was first described by Hager in **1885** ²⁾.

Bernhardt reported more extensively on the condition in **1895**, and 2 weeks later Roth published a paper in which he coined the term **meralgia**, from the Greek words meros for thigh and algos for pain ³⁾. and Bernhardt in **1895** ⁴⁾.

¹⁾

Bohmfolk GL, Story JL, Wissinger JP, et al. Bacterial Intracranial Aneurysm. J Neurosurg. 1978; 48:369–382

²⁾

Wilson SAK. Neurology, 2nd ed. Baltimore: Williams & Wilkins; 1955: 369.

³⁾

Roth: Meralgia Pareathetica, Berlin, S. Karger, 1895.

⁴⁾

Bernhardt Ueber isoliert Gibiete des Nervua cutanea femoris externua vorkommende paraeathealen, Neurot. Centraibl 242, 1895.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

<https://neurosurgerywiki.com/wiki/doku.php?id=1885>

Last update: **2024/06/07 02:58**

