

1917

1916-1918

The unilateral [suboccipital approach](#) was popularized by Woolsey (1903) and with great contributions by [Fedor Krause](#) (1905).

After several refinements and modifications through different dedicated neurosurgeons (Fish, House and Seiffert , Dandy's suboccipital approach (1917) with an ipsilateral suboccipital flap evolved to the [retrosigmoid transmeatal approach](#).

[Infundibulo tuberal syndrome](#) groups endocrine, metabolic, and behavioral disturbances caused by lesions involving the upper [neurohypophysis](#) (median eminence) and adjacent basal [hypothalamus](#) ([tuber cinereum](#)). It was originally described by Henri Claude and Jean Lhermitte in 1917, in a patient with a [craniopharyngioma](#) ¹⁾.

¹⁾

Castro-Dufourny I, Carrasco R, Prieto R, Barrios L, Pascual JM. The infundibulo-tuberal syndrome caused by craniopharyngiomas: clinicopathological evidence from an historical French cohort (1705-1973). Pituitary. 2014 Dec 20. [Epub ahead of print] PubMed PMID: 25527245.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

<https://neurosurgerywiki.com/wiki/doku.php?id=1917>

Last update: **2024/06/07 02:51**

