

The cerebral metabolic rate of oxygen consumption (**CMRO₂**) arises from **neurons** utilizing energy for two functions:

1) maintenance of cell integrity (**homeostasis**) which normally accounts for \approx 40% of energy consumption, and 2) conduction of electrical impulses. The occlusion of an artery produces a central core of ischemic tissue where the **CMRO₂** is not met. The oxygen deficiency precludes aerobic **glycolysis** and **oxidative phosphorylation**. **ATP** production declines and cell homeostasis cannot be maintained, and within minutes irreversible cell death occurs; a so-called **cerebral infarction**. Surrounding this central core is the **penumbra**, where collateral flow (usually through **leptomeningeal vessels**) provides marginal **oxygenation** which may impair cellular function without immediate irreversible damage. Cells in the penumbra may remain viable for hours.

Cerebral ischemia due to **cerebral vasospasm** is a feared complication in patients following **aneurysmal subarachnoid hemorrhage (SAH)**.

see **Posttraumatic cerebral infarction**.

see **Malignant middle cerebral artery infarction**.

Brain infarction results from a focal decrease in **cerebral blood flow**.

A **tandem occlusion** is an uncommon presentation of **acute ischemic stroke** that involves occlusion of the extracranial **internal carotid artery** (EICA) and concomitant occlusion of either the intracranial ICA and/or **middle cerebral artery**

From:
<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:
https://neurosurgerywiki.com/wiki/doku.php?id=acute_ischemic_stroke_etiology

Last update: **2025/04/29 20:23**

