

Aneurysmal subarachnoid hemorrhage risk factors

Hypertension

Oral contraceptives

Cigarette [smoking](#)

Following [cocaine](#) abuse

[Alcohol](#) consumption: controversial

Diurnal variations in [blood pressure](#)

[Pregnancy](#) and [parturition](#)

Slight increased risk during lumbar puncture and/or cerebral angiography in patient with cerebral aneurysm

Slight increased risk with advancing age

Conditions with an increased incidence of cerebral aneurysms

Left [ventricular assist device](#)-associated [subarachnoid hemorrhage](#) may be caused by [infectious intracranial aneurysms](#) when acute bloodstream infections are present ¹⁾.

¹⁾

Lee T, Buletko AB, Matthew J, Cho SM. Bloodstream infection is associated with subarachnoid hemorrhage and infectious intracranial aneurysm in left ventricular assist device. Perfusion. 2019 Jul 24;267659119858853. doi: 10.1177/0267659119858853. [Epub ahead of print] PubMed PMID: 31339450.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=aneurysmal_subarachnoid_hemorrhage_risk_factors

Last update: **2024/06/07 02:59**

