

Anterior Circulation Large Vessel Occlusion

- Association of hypoperfusion intensity ratio and cerebral blood volume Index with good outcome in patients transferred for thrombectomy
- Multicenter experience of the Monopoint reperfusion system in acute large vessel occlusion stroke thrombectomy
- Association Between Early Spontaneous Post-Thrombectomy Blood Pressure Reduction and Clinical Outcomes in Large Vessel Occlusion Stroke
- Predictors of futile recanalisation in patients with large infarct: a post-hoc analysis of the ANGEL-ASPECT trial
- Integrating perfusion imaging derived venous outflow and tissue-level collateral parameters in a comprehensive clinical model enhances prognostication in large vessel occlusion stroke
- Intra-arterial Tenecteplase for Acute Stroke After Successful Endovascular Therapy: The ANGEL-TNK Randomized Clinical Trial
- Intra-Arterial Tenecteplase After Successful Reperfusion in Large Vessel Occlusion Stroke: A Randomized Clinical Trial
- Petechial hemorrhage in mechanical thrombectomy for distal and medium-vessel occlusions: technical considerations and outcomes

Anterior Circulation LVO

- **Internal Carotid Artery (ICA)**
 - Cervical ICA (extracranial)
 - Intracranial ICA (supraclinoid segment)
- **Middle Cerebral Artery (MCA)**
 - M1 segment: from ICA bifurcation to main branches
 - M2 segment: opercular branches in Sylvian fissure
- **Anterior Cerebral Artery (ACA)** – less frequent
 - A1 segment

Prognosis

In a [retrospective cohort study](#) Asimos et al. from Atrium Health, Charlotte (Emergency Medicine, Neurosciences Institute, Quality Analytics, Radiology, Neurosurgery, Neurology) published in the [Interventional Neuroradiology Journal](#) to assess whether [hypoperfusion](#) intensity ratio (HIR) and [cerebral blood volume index](#) (CBVI) measured via [CT perfusion](#) at referring non-thrombectomy centers predict favorable 90-day outcomes post-transfer for [thrombectomy](#) in [anterior circulation large vessel occlusion](#) (ACLVO). CBVI—as a continuous measure and specifically > 0.7 —correlated with functional independence ($mRS \leq 2$) at 90 days both overall and in recanalized patients. In contrast, HIR thresholds and combined collateral scores were not predictive ¹⁾.

Critical Review

Strengths

Excellent [sample size](#) ($n = 497$), with high treatment prevalence (93% thrombectomy) supporting real-

world relevance.

Rigorous adjustment for confounders in multivariable models enhances validity.

Weaknesses

Retrospective and single-center design at a large referral system may limit external [generalizability](#).

No direct comparisons to other perfusion metrics like Tmax or ischemic core volumes—makes it difficult to situate CBVI within broader CTP prognostication tools.

HIR binary thresholds (0.4–0.6) may lack granularity; their univariate non-association could stem from arbitrary cutoffs rather than biological irrelevance.

Unclear reproducibility or inter-observer reliability of CBVI quantification across centers or software versions.

Statistical note

Adjusted OR of 1.73 for CBVI > 0.7 is clinically meaningful, but [confidence interval](#) (1.13–2.65) suggests moderate precision.

Final Verdict

Rating: 6.5 / 10

Takeaway for practitioners: CBVI from CT perfusion at referring centers could offer a pragmatic predictor of functional independence after thrombectomy, especially when > 0.7. However, its standalone prognostic value remains uncertain until validated prospectively and benchmarked against established perfusion metrics.

Bottom line: CBVI appears promising as a simple, transportable predictor of favorable outcome in ACLVO, but further multi-center, prospective validation and comparison studies are needed before adoption into routine transfer decision-making.

Categories: Retrospective Studies, Stroke Imaging, Thrombectomy Outcomes

Tags: CT perfusion, CBVI, HIR, collateral perfusion, stroke prognostication, thrombectomy, anterior circulation LVO

1)

Asimos AW, Yang H, Strong D, Teli KJ, Clemente JD, DeFilipp G, Bernard J, Stetler W, Parish JM, Hines A, Rhoten JB, Karamchandani RR. Association of [hypoperfusion intensity ratio](#) and [cerebral blood volume Index](#) with good [outcome](#) in patients transferred for [thrombectomy](#). Interv Neuroradiol. 2025 Jul 10:15910199251352046. doi: 10.1177/15910199251352046. Epub ahead of print. PMID: 40638076.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=anterior_circulation_large_vessel_occlusion&rev=1752180232

Last update: **2025/07/10 20:43**

