

# Anterior myelomeningocele treatment

see [Anterior sacral myelomeningocele treatment](#).

- [A "Smurf-Cap" head requiring total cranial vault reshaping. A novel syndromic presentation of craniofrontonasal dysplasia associated with spina bifida](#)
- [A case of spinal arteriovenous fistula complicated with congenital spinal lipoma and successfully treated with endovascular therapy](#)
- [Adult anterior sacral myelomeningocele associated with a didelphys uterus: A rare case report presentation](#)
- [Successful surgical management of anterior cervical meningomyelocele associated with Klippel-Feil deformity using anterior vertebral reconstruction: a case report](#)
- [Currarino syndrome with immature teratoma: A case report with review of literature](#)
- [The impact of shoes versus ankle-restricted orthoses on sit-to-stand kinematics and centre of mass trajectories in adults with myelomeningocele](#)
- [Treatment of an anterior cervicothoracic myelomeningocele together with spine deformity correction in a child: illustrative case](#)
- [Factors affecting clean intermittent catheterization compliance among children and adolescents with neurogenic bladder due to spina bifida and caudal regression syndrome](#)

Treatment for [anterior myelomeningocele](#) typically involves surgical repair. The goal of surgery is to protect the exposed spinal cord and nerves from damage and to prevent infection. The surgery may be performed shortly after birth, or it may be delayed until the child is several months old. In some cases, multiple surgeries may be needed to correct the defect and address associated complications.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

[https://neurosurgerywiki.com/wiki/doku.php?id=anterior\\_myelomeningocele\\_treatment](https://neurosurgerywiki.com/wiki/doku.php?id=anterior_myelomeningocele_treatment)

Last update: **2024/06/07 02:53**

