

Carvalho et al. investigated the impact of [c-Met](#), [HGF](#), [VEGFR2](#) expression and [microvessel density](#) (MVD) in Glioblastoma patients submitted to second-line chemotherapy with [bevacizumab](#). Immunohistochemical expression of c-Met, HGF, VEGFR2, and MVD was assessed in tumor specimens of Glioblastoma patients treated with bevacizumab, after progression under [temozolomide](#). Survival analysis was evaluated according to the expression of the aforementioned biomarkers. c-Met overexpression was associated with a time-to-progression (TTP) after bevacizumab of 3 months (95% CI, 1.5-4.5) compared with a TTP of 7 months (95% CI, 4.6-9.4) in patients with low or no expression of c-Met ($p = 0.05$). VEGFR2 expression was associated with a TTP after bevacizumab of 3 months (95% CI, 1.8-4.2) compared with a TTP of 7 months (95% CI, 5.7-8.3) in patients with no tumoral expression of VEGFR2 ($p = 0.009$). Concomitant c-Met/VEGFR2 overexpression was associated with worse overall survival (13 months) compared with concomitant c-Met/VEGFR2 negative expression (19 months; $p = 0.025$). This data support the hypothesis that c-Met and VEGFR2 overexpression have a role in the development of glioblastoma early resistance and might predict poorer responses to [anti-angiogenics](#) ¹⁾.

¹⁾

Carvalho B, Lopes JM, Silva R, Peixoto J, Leitão D, Soares P, Fernandes AC, Linhares P, Vaz R, Lima J. The role of [c-Met](#) and [VEGFR2](#) in [glioblastoma](#) resistance to [bevacizumab](#). Sci Rep. 2021 Mar 16;11(1):6067. doi: 10.1038/s41598-021-85385-1. PMID: 33727583.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

<https://neurosurgerywiki.com/wiki/doku.php?id=anti-angiogenics>

Last update: **2024/06/07 02:58**

