

Horizontal fractures of the atlas are uncommon fractures associated with instability of the [craniocervical junction](#). Most commonly associated with high-speed motor vehicle accidents, these fractures need to be identified and treated appropriately. Due to its relatively benign presentation on bony imaging, magnetic resonance imaging to look for ligamentous instability is important. We present two such cases which were managed by [occipitocervical fusion](#) at our institute ¹⁾.

This is the first report of traumatic combined vertical atlanto-occipital dislocation (AOD) and atlanto-axial dislocation (AAD) with 2-part fracture of the atlas.

Patient concerns: The first case was of a 31-year-old woman admitted to the emergency room comatose after a traffic accident. The second case was of a 21-year-old woman admitted to the emergency room comatose after a fall.

Diagnoses: Traumatic combined vertical AOD and AAD with 2-part fractures of the atlas was diagnosed using plain radiography, 2-dimensional computed tomography, and/or magnetic resonance imaging of the cervical spine.

Intervention: The first patient received immediate intubation and cardiopulmonary resuscitation in the emergency room. The second patient also received immediate intubation in the emergency room. After her vitals stabilized, she underwent occipitocervical fusion with instrumentation.

Outcomes: The first patient died 2 days after the accident. The second patient remained quadriplegic in a ventilatory-dependent state at 1 year after surgery. She continues to receive comprehensive rehabilitation ²⁾.

A 4-year-old child in whom a [penetrating trauma](#) to an immature [atlas](#) led to an unusual disjunction of the posterior [synchondrosis](#) with [atlas fracture](#) displacement of the posterior “hemiarch” that plunged into the [dura](#), resulting in a [cerebrospinal fluid fistula](#).

Praneeth et al. discussed the possible mechanism and considerations in the [management](#) of this unique presentation. Such an atypical fracture pattern involving the posterior hemi ring of the pediatric atlas is previously unknown ³⁾.

¹⁾

Brown J, Hegde A. Horizontal fracture of the atlas - A rare but unstable C1 fracture. J Craniovertebr Junction Spine. 2020 Oct-Dec;11(4):338-341. doi: 10.4103/jcvjs.JCVJS_143_20. Epub 2020 Nov 26. PMID: 33824565; PMCID: PMC8019112.

²⁾

Park JB, Chang DG, Kim WJ, Kim ES. Traumatic combined vertical atlanto-occipital and atlanto-axial dislocations with 2-part fracture of the atlas: Two case reports. Medicine (Baltimore). 2019 Nov;98(44):e17776. doi: 10.1097/MD.00000000000017776. PMID: 31689843; PMCID: PMC6946329.

³⁾

Praneeth K, Karthigeyan M, Salunke P, Ray N. Synchondral Fracture of the Posterior “Hemiarch” of Pediatric Atlas with Cerebrospinal Fluid Fistula following a Penetrating Neck Injury. Pediatr Neurosurg. 2019 Oct 10:1-4. doi: 10.1159/000503109. [Epub ahead of print] PubMed PMID: 31600753.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=atlas_fracture_case_reports

Last update: **2024/06/07 02:48**

