

# Claw sign

Modic type 1 degenerative signal changes can mimic/suggest infection, leading to additional costly and sometimes invasive investigations.

A retrospective study analyzes the utility and accuracy of a novel, diffusion-weighted “claw sign” for distinguishing symptomatic type 1 degeneration from vertebral diskitis/osteomyelitis.



In patients with type 1 signal changes of the vertebral disk space, a claw sign is highly suggestive of degeneration and its absence strongly suggests diskitis/osteomyelitis <sup>1)</sup>.

<sup>1)</sup>

Patel KB, Poplawski MM, Pawha PS, Naidich TP, Tanenbaum LN. Diffusion-weighted MRI “claw sign” improves differentiation of infectious from degenerative modic type 1 signal changes of the spine. AJNR Am J Neuroradiol. 2014 Aug;35(8):1647-52. doi: 10.3174/ajnr.A3948. Epub 2014 Apr 17. PubMed PMID: 24742801.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**



Permanent link:

[https://neurosurgerywiki.com/wiki/doku.php?id=claw\\_sign](https://neurosurgerywiki.com/wiki/doku.php?id=claw_sign)

Last update: **2025/04/29 20:23**