

- Borders: 1. [Optic Nerve](#)
- 2. [Oculomotor Nerve](#)
- 3. [Tentorial Edge](#)
- Contents: 1. [Anterior Clinoid Process](#)
- 2. [Clinoidal ICA](#)

The clinoidal triangle can be exposed via the endoscopic supraorbital extradural approach. Following a [supraorbital craniotomy](#) and the [drilling](#) of the [sphenoidal ridge](#), the [orbital roof](#) and the greater and [lesser sphenoid wings](#), the floor of the [middle cranial fossa](#) can be exposed by lifting the temporal dura slightly. At this point, the [anterior clinoid process](#) can be removed to allow the clinoidal triangle to be visualized ¹⁾.

1)

Komatsu F, Komatsu M, Inoue T, Tschabitscher M. Endoscopic supraorbital extradural approach to the cavernous sinus: a cadaver study. J Neurosurg. 2011 May;114(5):1331-7. doi: 10.3171/2010.10.JNS101242. Epub 2010 Dec 10. PMID: 21142751.

From:
<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**



Permanent link:
https://neurosurgerywiki.com/wiki/doku.php?id=clinoidal_triangle

Last update: **2024/06/07 02:51**