

Convexity meningioma

J.Sales-Llopis

Neurosurgery Service, *General University Hospital Alicante*, Spain.

Latest Pubmed Related Articles

- [A Case Report Depicting a Rare Neurosurgical Disease: Aggressive Meningiomatosis](#)
- [Perioperative Seizures and Quality of Life in Falx and Convexity Meningiomas: Key Factors of Patient Outcomes](#)
- [Comprehensive analysis of the distribution of dural artery-derived tumor-feeding arteries in intracranial meningiomas](#)
- [Intracranial meningioma with intratumoral hemorrhage: A systematic review of associated features and outcomes](#)
- [En plaque convexity hyperostotic meningioma: 69 cases from a single center](#)
- [Chronic calcified subdural hematoma masquerading as hemorrhagic extra-axial mass: A case report and brief review of the literature](#)
- [Hyperostotic meningiomas in children. A case-based update](#)
- [Emergency Surgical Management of Meningiomas: Factors Affecting Early Outcomes and Complications](#)

Intracranial meningioma that grow on the surface of the brain directly under the skull.

Classification

[Convexity meningioma classification.](#)

Epidemiology

[Convexity meningioma epidemiology.](#)

Clinical features

[Convexity meningioma clinical features.](#)

Diagnosis

[Convexity meningioma diagnosis.](#)

Differential diagnosis

[Convexity meningioma differential diagnosis.](#)

Treatment

[Convexity meningioma treatment.](#)

Outcome

[Convexity meningioma outcome.](#)

Complications

[Convexity meningioma complications.](#)

Case series

[Convexity meningioma case series.](#)

Case reports

[Convexity meningioma case reports.](#)

Case reports from the HGUA

Q11705

64-year-old male


Medical History: Allergy to [Nonsteroidal anti-inflammatory drugs](#), [ex-smoker](#) (30 years), [bronchopathy \(asthma\)](#), [dyslipidemia](#), mixed [hearing loss](#), [thrombocytopenia](#) under investigation.

Current Medication: Atorvastatin 30 mg, Mebhydrolin 150 mg, Salbutamol inhaler 100 mcg/1 puff, Paracetamol 1000 mg, Azithromycin 500 mg, Tamsulosin/Solifenacin 0.4/6 mg.

The patient presented with a history of malaise, headache, nausea, vomiting, confusion, and disorientation over the past few weeks. Admission was prompted by a cerebral loss of consciousness.

The patient was conscious and oriented, without neurological focal deficits.

Complementary Examination:




Solid mass in the right frontal lobe, approximately 144 x 39.7 x 31.5 mm in size, causing significant perilesional edema. Suspected to be an atypical meningioma or hemangiopericytoma.

[Atypical meningioma](#) or [hemangiopericytoma](#) causing significant [perilesional edema](#).

CT-Thorax and Abdomen-Pelvis: Unremarkable, except for a 6 mm hepatic cyst in segment 8.

This case report highlights the challenging presentation of an atypical meningioma in an elderly male, emphasizing the need for prompt diagnosis and intervention. The upcoming surgery aims to address the cerebral mass and associated complications, with careful consideration of the patient's overall health and ongoing medical management. Further updates and outcomes will be documented following the intervention.



From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=convexity_meningioma

Last update: **2025/04/29 20:30**

