

Cord signal change (CSC)

An important finding of intrinsic pathology is the presence of increased signal in the cervical spinal cord on [T2 weighted image](#), or [cord signal change](#) (CSC). This was first noted in the late 1980s and early 1990s ^{1) 2) 3)}.

CSC is thought to represent pathological changes in the [spinal cord](#) detectable with histology that occur as a result of chronic compression ⁴⁾. Many surgeons consider the presence of CSC indicative of advanced [cervical spondylotic myelopathy](#) (CSM), and a strong indication for surgical intervention. However, the significance of CSC for disease severity and prognosis remains controversial. Although several authors have shown that CSC is a reliable indicator of surgical outcome, ^{5) 6) 7) 8) 9)} others have not ^{10) 11) 12) 13)}.

Results suggest that low intensity signal on preoperative [T1 weighted image](#), but not T2-WIs correlated with poor postoperative [neurological outcome](#). Furthermore, decreased signal intensity on postoperative T1-WIs and increased signal intensity on postoperative T2-WIs are predictors of poor neurologic outcome.

Despite the numerous studies examining the relationship of CSC with outcomes of both surgical and nonsurgical treatment of CSM, there have not been any studies examining specifically the relationship of physical examination findings with CSC.

CSC visualized on MRI correlates poorly with the upper extremity reflex examination in patients with cervical myelopathy. Of the pathological reflexes, [Hoffmann's reflex](#) has the strongest association with CSC, but still was only positive in 67% of cases. More sensitive clinical measures need to be developed to more accurately associate CSC detected on MRI to the clinical severity of cervical spondylotic myelopathy. ¹⁴⁾

References

^{1) 5)}

Mehalic TF, Pezzuti RT, Applebaum BI. Magnetic resonance imaging and cervical spondylotic myelopathy. *Neurosurgery* 1990;26:217-26; discussion 26-7.

²⁾

Takahashi M, Sakamoto Y, Miyawaki M, et al. Increased MR signal intensity secondary to chronic cervical cord compression. *Neuroradiology* 1987;29:550-6.

³⁾

Takahashi M, Yamashita Y, Sakamoto Y, et al. Chronic cervical cord compression: clinical significance of increased signal intensity on MR images. *Radiology* 1989;173:219-24.

⁴⁾

Ohshio I, Hatayama A, Kaneda K, et al. Correlation between histopathologic features and magnetic resonance images of spinal cord lesions. *Spine (Phila Pa 1976)* 1993;18:1140-9.

⁶⁾

Al-Mefty O, Harkey LH, Middleton TH, et al. Myelopathic cervical spondylotic lesions demonstrated by magnetic resonance imaging. *J Neurosurg* 1988;68:217-22.

⁷⁾

Matsuda Y, Miyazaki K, Tada K, et al. Increased MR signal intensity due to cervical myelopathy. Analysis of 29 surgical cases. *J Neurosurg* 1991;74:887-92.

⁸⁾

Okada Y, Ikata T, Yamada H, et al. Magnetic resonance imaging study on the results of surgery for cervical compression myelopathy. *Spine (Phila Pa 1976)* 1993;18:2024-9.

9)

Wada E, Yonenobu K, Suzuki S, et al. Can intramedullary signal change on magnetic resonance imaging predict surgical outcome in cervical spondylotic myelopathy? Spine (Phila Pa 1976) 1999;24:455-61; discussion 62.

10)

Morio Y, Yamamoto K, Kuranobu K, et al. Does increased signal intensity of the spinal cord on MR images due to cervical myelopathy predict prognosis? Arch Orthop Trauma Surg 1994;113:254-9.

11)

Naderi S, Ozgen S, Pamir MN, et al. Cervical spondylotic myelopathy: surgical results and factors affecting prognosis. Neurosurgery 1998;43:43-9; discussion 9-50.

12)

Nakamura M, Fujimura Y. Magnetic resonance imaging of the spinal cord in cervical ossification of the posterior longitudinal ligament. Can it predict surgical outcome? Spine (Phila Pa 1976) 1998;23:38-40.

13)

Yone K, Sakou T, Yanase M, et al. Preoperative and postoperative magnetic resonance image evaluations of the spinal cord in cervical myelopathy. Spine (Phila Pa 1976) 1992;17:S388-92

14)

Nemani VM, Kim HJ, Piyaskulkaew C, Nguyen JT, Riew KD. Correlation of cord signal change with physical examination findings in patients with cervical myelopathy. Spine (Phila Pa 1976). 2015 Jan 1;40(1):6-10. doi: 10.1097/BRS.0000000000000659. PubMed PMID: 25341986.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=cord_signal_changeLast update: **2025/04/29 20:27**