

Cytokine receptor superfamily

The [cytokine receptor](#) superfamily is a large group of cell surface receptors that mediate the actions of various [cytokines](#). Cytokines are [signaling molecules](#) that play crucial roles in regulating immune responses, inflammation, cell growth, differentiation, and various physiological processes. Cytokine receptors are classified into several families based on their structural and functional characteristics, including the Class I, Class II, and Class III cytokine receptor families. These receptors are key components of the immune system and have wide-ranging effects on cell function and development.

Here are some important characteristics and examples of cytokine receptor families within the cytokine receptor superfamily:

Class I Cytokine Receptors

These receptors have a single transmembrane domain and are associated with the activation of the Janus kinase-signal transducer and activator of the transcription (JAK-STAT) signaling pathway. Examples of Class I cytokine receptors include receptors for interleukins (e.g., IL-2, IL-6, IL-10), interferons (e.g., IFN- α , IFN- β), and granulocyte colony-stimulating factor (G-CSF).


Class II Cytokine Receptors: Class II cytokine receptors also have a single transmembrane domain but signal through different pathways. Examples include the receptors for interferon- γ (IFN- γ) and the interleukin-10 receptor.

Class III Cytokine Receptors: These receptors have two or more extracellular fibronectin type III domains and play a role in cell proliferation and differentiation. An example is the erythropoietin receptor, which is involved in the regulation of red blood cell production.

Homodimeric and Heterodimeric Receptors: Cytokine receptors can form homodimers (two identical receptor subunits) or heterodimers (two different receptor subunits) upon cytokine binding, allowing for diverse responses to various cytokines.

Cytokine receptor activation triggers intracellular signaling cascades that regulate gene expression, control immune cell functions, and coordinate various biological responses. Dysregulation of cytokine signaling or mutations in cytokine receptors can lead to immune-related disorders, autoimmune diseases, and other health conditions.

The cytokine receptor superfamily is an essential part of the immune and inflammatory responses, and it is a target for therapeutic interventions in diseases involving immune system dysregulation.



From:
<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:
https://neurosurgerywiki.com/wiki/doku.php?id=cytokine_receptor_superfamily

Last update: **2024/06/07 02:58**

