

Diffuse infiltrative gliomas are by far the most common primary brain tumors in adults, esp. its most malignant form, [glioblastoma multiforme](#) (Glioblastoma) ¹⁾.

Diffuse gliomas are the second most common [central nervous system tumors](#), behind meningiomas, and account for roughly 80% of primary malignant brain tumors ^{2) 3) 4)}.

¹⁾
CBTRUS (2006) CBTRUS statistical report. Primary brain tumors in the United States, 1998–2002. Central Brain Tumor Registry of the United States. <http://www.cbtrus.org>

²⁾
Bondy ML, Scheurer ME, Malmer B, et al. Brain tumor epidemiology: consensus from the brain tumor epidemiology consortium (BTEC). Cancer 2008;113:1953–68

³⁾
Jansen M, Yip S, Louis DN. Molecular pathology in adult gliomas: diagnostic, prognostic and predictive markers. Lancet Neurol 2010;9:717–26

⁴⁾
Theeler BJ, Yung WK, Fuller GN, et al. Moving toward molecular classification of diffuse gliomas in adults. Neurology 2012;79:1917–26

From:
<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:
https://neurosurgerywiki.com/wiki/doku.php?id=diffuse_glioma_epidemiology

Last update: **2024/06/07 02:52**

