


Disc height index

DHI ([disc height](#) index) is measured by ab/bc and FHI (foraminal height index) is measured by de/bc . Ab indicates the disc height, bc indicates the vertebrae height and de indicates the longest distance between the upper and lower vertebral pedicle. Ac represents the line labeling the centers. The center of the vertebral body is marked by the crossing point of two diagonal lines



Moriguchi et al., demonstrated that [riboflavin](#) crosslinked high-density [collagen gels](#) (HDC) can facilitate annular repair in vivo.

42 rats, tail disc punctured with an 18-gauge needle, were divided into 3 groups: untreated (n=6), injected with crosslinked HDC (n=18), and injected with [Annulus Fibrosus cell](#)-laden crosslinked HDC (n=18). Ovine AF cells were mixed with HDC gels prior to injection. X-rays and MRIs were conducted over 5 weeks, determining [disc height index](#) (DHI), [nucleus pulposus](#) (NP) size, and hydration. Histological assessments evaluated the viability of implanted cells and degree of annular repair.

Although average DHIs of both HDC gel groups were higher than those of the puncture control group at 5 weeks, the retention of disc height, NP size and hydration at 1 and 5 weeks was significant for the

cellular group compared to the punctured, and at 5 weeks to the acellular group. Histological assessment indicated that AF cell-laden HDC gels have accelerated reparative sealing compared to acellular HDC gels.

AF cell-laden HDC gels have the ability of better repairing annular defects than acellular gels after needle puncture.

This project addresses the compelling demand of a sufficient treatment strategy for [degenerative disc disease](#) (DDD) perpetuated by annulus fibrosus (AF) injury, a major cause of morbidity and burden to health care systems.

The study is designed to answer the question of whether injectable, photo-crosslinked, high density collagen gels can seal defects in the annulus fibrosus of rats and prevent [disc degeneration](#). Furthermore, we investigated whether the healing of AF defects will be enhanced by the delivery of AF cells (fibrochondrocytes) to these defects. The use of cell-laden collagen gels in spine surgery holds promise for a wide array of applications, from current [discectomy](#) procedures to future [nucleus pulposus](#) reparative therapies ¹⁾.

¹⁾

Moriguchi Y, Borde B, Berlin C, Wipplinger C, Sloan SR, Kirnaz S, Pennicooke B, Navarro-Ramirez R, Khair T, Grunert P, Kim E, Bonassar L, Härtl R. In Vivo Annular Repair using High-Density Collagen Gel Seeded with Annulus Fibrosus Cells. *Acta Biomater*. 2018 Jul 4. pii: S1742-7061(18)30402-1. doi: 10.1016/j.actbio.2018.07.008. [Epub ahead of print] PubMed PMID: 29981494.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=disc_height_index

Last update: **2024/06/07 02:58**

