Escala de TAC de Rotterdam

Esta escala se basa en las anomalías TAC tras un TCE

Los elementos de puntuación son:

Cisternas basales

0: normales

1: comprimidas

2: ausentes

Desviación de línea media

0: no hay cambios o <5 mm

1: > 5 mm

Lesión epidural

0: presente

1: ausente

Hemorragia intraventricular o subaracnoidea

0: ausente

1: presente

Instrucciones para su uso

La puntuación final es la suma de los elementos de puntuación + 1.

La mortalidad a los 6 meses es:

Resultado 1: 0%

Resultado 2: 7%

Resultado 3: 16%

Resultado 4: 26%

Puntuación 5: 53%

Resultado 6: 61% [1].

Bibliografía

1)

Maas AIR, Hukkelhoven CWPM, Marshall LF, Steyerberg EW. Prediction of outcome in traumatic brain injury with computed tomographic characteristics: a comparison between the computed tomographic classification and combinations of computed tomographic predictors. Neurosurgery. 2005 Dec.;57(6):1173-82

From:
https://neurosurgerywiki.com/wiki/ - **Neurosurgery Wiki**

Permanent link:
https://neurosurgerywiki.com/wiki/doku.php?id=escala_tac_de_rotterdam

Last update: **2025/05/13 02:26**