








Evoked potential waveforms

Test	Figure	Possible generators
BAER		CM
		cochlear microphonic
		P ₁
		distal VIII nerve,
		P ₂
		proximal VIII or cochlear nucleus,
UE SSEP		P ₃
		lower pons (? superior olivary complex),
		P ₄
		mid-upper pons,
		P ₅
		upper pons or inferior colliculus
LE SSEP		N ₉
		(on F _z -E _p where E _p is Erb's point) AKA EP: entry of volley into distal brachial plexus,
		N ₁₁
		(on C _{V7} -F _{Pz}): root entry zone (cervical region),
		N ₁₃
		cervicomedullary junction (recorded from C ₂),
PR VER		N ₁₉
		primary sensory cortex,
		P ₂₂
		(early) motor cortex,
		P ₂₂
		(late) IPSP "reaction" to N ₁₈
PR VER		P ₂₂
		(on L ₅ -T ₁₂): lumbo-sacral plexus,
		P ₄₀
		(on C _z -F _{Pz}): sensory cortex (analogous to N ₁₈ in UE SSEP, reversed in polarity for ? reason),
		N ₂₇
		(on C _{V7} -F _{Pz}): ? dorsal column nucleus
PR VER		N ₃₀
PR VER		P ₁₀₀
		(striate & pre-striate occipital cortex, with contributions from thalamocortical volleys)

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=evoked_potential_waveforms

Last update: **2024/06/07 02:58**

