

Excitability

Excitability refers to the ability of neurons and other cells to respond to stimuli by generating electrical signals called action potentials. It is a fundamental property of neurons that allows them to communicate with each other and with other types of cells, such as muscle cells, to produce coordinated responses in the body.

Key Aspects of Neuronal Excitability: Resting Membrane Potential:

Definition: The resting membrane potential is the difference in electric charge across the cell membrane when a neuron is not actively firing. It is typically around -70 mV in neurons, with the inside of the cell being more negative compared to the outside. **Ionic Basis:** The resting membrane potential is maintained by the uneven distribution of ions (such as sodium, potassium, and chloride) across the cell membrane, and the action of ion pumps, particularly the sodium-potassium pump (Na⁺/K⁺ ATPase). **Action Potential:**

Definition: An action potential is a rapid, temporary change in the membrane potential that travels along the axon of a neuron. It is the primary way that neurons transmit information. **Phases:** **Depolarization:** Triggered by a stimulus, voltage-gated sodium channels open, allowing Na⁺ ions to rush into the cell, causing the membrane potential to become more positive. **Repolarization:** After depolarization, voltage-gated potassium channels open, allowing K⁺ ions to flow out of the cell, restoring the membrane potential to a negative value. **Hyperpolarization:** The membrane potential may temporarily become more negative than the resting potential due to the continued efflux of K⁺ ions before stabilizing again at the resting potential. **Threshold:**

Definition: The threshold is the critical level of depolarization that must be reached for an action potential to be initiated. If the membrane potential does not reach this threshold, the neuron will not fire. **All-or-None Principle:** Once the threshold is reached, an action potential will occur in an all-or-none fashion, meaning it either happens fully or not at all. **Ionic Channels and Excitability:**

Voltage-Gated Channels: Neuronal excitability is heavily dependent on the function of voltage-gated ion channels, such as sodium, potassium, and calcium channels. These channels open or close in response to changes in membrane potential, controlling the flow of ions and the generation of action potentials. **Modulation:** Excitability can be modulated by various factors, including neurotransmitters, neuromodulators, and drugs that alter the function of these ion channels. **Synaptic Input:**

Excitatory Postsynaptic Potentials (EPSPs): Synaptic input that depolarizes the membrane potential, bringing it closer to the threshold for an action potential, increases excitability. This is typically mediated by neurotransmitters like glutamate that open cation channels (e.g., Na⁺ channels). **Inhibitory Postsynaptic Potentials (IPSPs):** Synaptic input that hyperpolarizes the membrane potential, moving it further from the threshold, decreases excitability. This is typically mediated by neurotransmitters like GABA that open anion channels (e.g., Cl⁻ channels). **Intrinsic and Extrinsic Factors Influencing Excitability:**

Intrinsic Factors: These include the properties of the neuron's membrane, the density and types of ion channels present, and the neuron's overall health and metabolic state. **Extrinsic Factors:** These include the influence of other neurons, hormones, and external stimuli. For instance, neuromodulators like serotonin or dopamine can change a neuron's excitability by affecting ion channel activity. **Clinical Relevance:**

Epilepsy: Abnormal increases in neuronal excitability can lead to conditions like epilepsy, where excessive, synchronized firing of neurons results in seizures. Anxiety and Mood Disorders: Alterations in the excitability of neurons in certain brain regions can contribute to anxiety, depression, and other psychiatric conditions. Neurodegenerative Diseases: Changes in neuronal excitability are also implicated in neurodegenerative diseases like Alzheimer's and Parkinson's, where disrupted signaling contributes to cognitive and motor dysfunction. In summary, excitability is a crucial property of neurons that underlies their ability to generate action potentials and communicate within the nervous system. It is finely regulated by the interaction of various ionic channels, synaptic inputs, and both intrinsic and extrinsic factors. Disruptions in excitability can lead to a range of neurological and psychiatric disorders.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

<https://neurosurgerywiki.com/wiki/doku.php?id=excitability>

Last update: **2025/05/13 02:01**

