

Fourth ventricle choroid plexus papilloma

In childhood, [choroid plexus papilloma](#) CPPs are usually found in the [lateral ventricles](#) and less commonly in the [posterior fossa](#). On the contrary, in [adults](#), the majority of these [tumors](#) are found in the [fourth ventricle](#) and its [lateral recesses](#) ¹⁾.

A case of 72-year-old man with posterior fossa lesion operated with provisional diagnosis of metastasis, but turn out to be CPP ²⁾.

Differential diagnosis

Medulloblastoma.

[Atypical teratoid/rhabdoid tumor](#) (AT/RT).

Ependymoma.

```
<html><iframe width="560" height="315" src="https://www.youtube.com/embed/Xbd2MwC7Wu4"
frameborder="0" allow="autoplay; encrypted-media" allowfullscreen></iframe></html>
```

¹⁾

Lee SH, Park BJ, Kim EJ, Lim YJ. Atypical choroid plexus papilloma in an adult. J Korean Neurosurg Soc. 2009 Jul;46(1):74-6. doi: 10.3340/jkns.2009.46.1.74. Epub 2009 Jul 31. PubMed PMID: 19707500; PubMed Central PMCID: PMC2729831.

²⁾

Umredkar AA, Chhabra R, Bal A, Das A. Choroid plexus papilloma of the fourth ventricle in a septuagenarian. J Neurosci Rural Pract. 2012 Sep;3(3):402-4. doi: 10.4103/0976-3147.102643. PubMed PMID: 23189015; PubMed Central PMCID: PMC3505354.

From:

<https://neurosurgerywiki.com/wiki/> - Neurosurgery Wiki

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=fourth_ventricle_choroid_plexus_papilloma

Last update: **2024/06/07 02:57**

