

Case series

From January 2006 to June 2015, 157 consecutive patients (98 females; mean age = 48.3 years) who suffered from meningiomas and were preoperatively referred for [embolization](#) were retrospectively analyzed. [Glubran](#) (n = 40) and PVA (n = 55) were used to devascularize tumors. Sixty-two patients were not embolized because of dangerous anastomosis or other tumor characteristics. Intraoperative blood loss, intraoperative time, degree of angiographic devascularization and embolization-related complications were analyzed.

The intraoperative blood loss and operative time were significantly lower in the Glubran-embolized versus non-embolized group. Furthermore, Glubran embolization significantly reduced intraoperative blood loss and operative time for meningiomas that received their primary blood supply from the ICA and/or VA compared with PVA embolization.

Preoperative meningioma embolization with Glubran decreases intraoperative blood loss and operative time. Furthermore, embolization with Glubran produces more effective devascularization compared with PVA for meningiomas supplied by the ICA and/or VA. Thus, Glubran may represent a better embolic agent for this meningioma subtype.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

<https://neurosurgerywiki.com/wiki/doku.php?id=glubran>

Last update: **2025/04/29 20:24**

