

Hemifacial spasm epidemiology

- [Identifying associated comorbidities in the development of trigeminal neuralgia: A propensity-matched analysis of the National Inpatient Sample](#)
- [Risk factors and Occurrence of Small vessel disease in Acute sensorineural hearing Loss In the Elderly: protocol for a multicentre cross-sectional study](#)
- [Which surgical technique has a superior clinical outcome in microvascular decompression? a systematic review and meta-analysis study of transposition versus interposition for trigeminal neuralgia and hemifacial spasm](#)
- [Fully Endoscopic Microvascular Decompression for Neurovascular Compression Syndrome: Early Outcomes and Technical Note](#)
- [Arterial Hypertension After Macrovascular Decompression of Vertebrobasilar Dolichoectasia: Rethinking the Cause-Effect Relationship](#)
- [Increasing Incidence of Facial Nerve Disorders in the United States from 2007 to 2022](#)
- [Efficacy and safety of reoperation for residual or recurrent hemifacial spasm after microvascular decompression: a systematic review and meta-analysis](#)
- [Optimized microvascular decompression surgery for improving the results of hemifacial spasm: an analysis of reoperations](#)

The incidence of hemifacial spasm is approximately 0.8 per 100,000 persons.

This disorder occurs in both men and women, although it affects middle-aged or elderly women more frequently.

Hemifacial spasm is much more common in some Asian populations.

From:
<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:
https://neurosurgerywiki.com/wiki/doku.php?id=hemifacial_spasm_epidemiology

Last update: **2024/06/07 02:57**

