

Herpes simplex virus

Infections are categorized based on the part of the body infected. Oral [herpes](#) involves the face or mouth. It may result in small blisters in groups often called cold sores or fever blisters or may just cause a sore throat.

Genital herpes, often simply known as herpes, may have minimal symptoms or form blisters that break open and result in small ulcers. These typically heal over two to four weeks. Tingling or shooting pains may occur before the blisters appear. Herpes cycles between periods of active disease followed by periods without symptoms. The first episode is often more severe and may be associated with fever, muscle pains, swollen lymph nodes and headaches. Over time, episodes of active disease decrease in frequency and severity.

Other disorders caused by herpes simplex include: herpetic whitlow when it involves the fingers, herpes of the eye, herpes infection of the brain, and neonatal herpes when it affects a newborn, among others.

see [Herpes simplex virus type 1](#)

see [Oncolytic herpes simplex virus](#).

Herpes simplex encephalitis

[Herpes simplex encephalitis](#).

Herpes simplex virus hydrocephalus

see [Herpes simplex virus hydrocephalus](#).

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=herpes_simplex_virus

Last update: **2025/04/29 20:25**

