

Horizontal gaze palsy

The maintenance of horizontal gaze is an essential function of upright posture and global sagittal spinal alignment. Horizontal gaze is classically measured by the **chin brow vertical angle** (CBVA), which is not readily measured on most lateral spine radiographs.

Horizontal **gaze palsy** may be caused by lesions in the cerebral hemispheres, which cause paresis of **gaze** away from the side of the lesion, or from **brainstem lesions**, which, if they occur below the crossing of the fibers from the **frontal eye fields** in the caudal midbrain, will cause weakness of gaze toward the side of the lesion. Another way to remember this is that patients with hemisphere lesions look toward their lesion, while patients with pontine gaze palsies look away from their lesions. Note that patients with gaze palsy still have conjugate eye movements and therefore do not complain of diplopia.

The human Robo gene acts as a receptor for a midline repulsive cue. When Robo is mutated, the longitudinal tract formation is disrupted and therefore normal neuronal connections cannot form. This leads to the reduced hindbrain volume and scoliosis, which are common symptoms of horizontal gaze palsy.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=horizontal_gaze_palsy

Last update: **2025/04/29 20:22**

