

Hypodense

Hypodense [lesions](#) identified on [Computerized tomography](#) are often assumed to indicate [ischemia](#).

see also [Computed Tomography for chronic subdural hematoma](#).



Doubtful hypodense left [occipital](#) intraparenchymal area, probably nonspecific, unable to rule out hemorrhagic parenchymal [brain contusion](#). Hospital observation and possibly CT control are recommended.

[Type 2 diabetes](#). [Atrial fibrillation](#), [Obesity](#), [chronic respiratory failure](#)

[Smoker](#) of 1-2 packs/day and severe [alcohol habit](#) Usual treatment -[MICARDIS PLUS 40/12.5MG](#) 1 tablet every 24 hours -[Dapagliflozin](#) and [metformin](#) combination 5MG/1000MG 1 tablet every 12 hours -[OMEPRazole](#) 20 MG 1 tab every 24h -[ELIQUIS](#) 5MG

see [Fall-Related Traumatic Brain Injury](#)

He [fall](#) the day before, he comments [loss of consciousness](#) lasting seconds, comments that he feels ill, referring to a sensation of [dizziness](#) with turning of objects, he refers to a sensation of relative [dyspnea](#), no vomiting, he refers to having drunk [alcohol](#)

[Contusion](#) in the [occipital region](#)

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

<https://neurosurgerywiki.com/wiki/doku.php?id=hypodense>

Last update: **2024/06/07 02:54**

