

Idiopathic intracranial hypertension surgery indications

- [Fulminant Idiopathic Intracranial Hypertension-Clinical Characteristics and Continuous Drainage as a Novel Treatment Approach](#)
- [Clinical indications and patient outcomes of intracranial venous sinus stenting beyond overt idiopathic intracranial hypertension: a scoping review](#)
- [Ventriculo-pleural shunt --- A second line option in the management of complex hydrocephalus](#)
- [Venous sinus stenting for management of spontaneous skull-base CSF leaks: A systematic review and meta-analysis](#)
- [Fibrin-coated collagen fleece "sandwich" closure technique for cranial and spinal dural reconstruction and closure](#)
- [The role of acetazolamide in critical care and emergency medicine](#)
- [The use of balloon guide catheters during venous sinus stenting: A case series](#)
- [Outcomes of Pulsatile Tinnitus After Cerebral Venous Sinus Stenting: Systematic Review and Pooled Analysis of 616 Patients](#)

Surgical intervention for idiopathic intracranial hypertension (IIH) is generally considered when medical management and other conservative treatments have failed or are insufficient. Here are the key indications and types of surgical interventions for IIH:

Indications for Surgery:

1. Persistent or Severe Symptoms:

1. When patients continue to experience severe headaches, visual disturbances, or other debilitating symptoms despite optimal medical treatment.

2. Visual Impairment:

1. If IIH is causing progressive vision loss or significant papilledema that threatens vision, and medical treatments are not effective in stabilizing or improving the condition.

3. Failure of Medical Management:

1. If patients do not respond adequately to medications like acetazolamide or other conservative treatments, surgical options may be considered.

4. Recurrent or Severe Papilledema:

1. In cases where papilledema (swelling of the optic nerve) is severe or recurrent despite treatment, and there is a risk of permanent vision damage.

5. Chronic Intracranial Pressure:

1. When there is a sustained increase in intracranial pressure that is not adequately controlled by medications or lifestyle changes.

From:
<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:
https://neurosurgerywiki.com/wiki/doku.php?id=idiopathic_intracranial_hypertension_surgery_indications

Last update: **2024/09/09 14:33**

