

iNPH scale

Examples of reported scales are the Black, Stein and Langfitt, Boon, Mori, and Krauss scales.

The iNPH scale developed by Hellström et al. measures symptoms and signs in 4 different domains: gait, cognition, continence, and balance ¹⁾.

Gait disorders

Bedridden or can not walk 5

You can walk with support 4

Walk unaided but unstable with frequent falls 3

stable but abnormal gait 2

Average 1

cognitive impairment

vegetative 5

severe dementia 4

serious changes in memory and behavior 3

Memory impairment as patient or family 2

Mild cognitive problems 1

sphincters

urinary and fecal incontinence 5

Continuous urinary incontinence 4

urinary incontinence sporadic 3

urinary urgency 2

normal 1

¹⁾

Hellström P, Klinge P, Tans J, Wikkelsø C. A new scale for assessment of severity and outcome in iNPH. *Acta Neurol Scand.* 2012 Oct;126(4):229-37. doi: 10.1111/j.1600-0404.2012.01677.x. Epub 2012 May 16. PubMed PMID: 22587624.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=inph_scale

Last update: **2024/06/07 02:57**

