

Intracranial aneurysm risk factors

- [Safety and Effect of Flow Diverters in the Management of Large and Giant Unruptured Intracranial Aneurysms](#)
- [Elevated Expression of TGFB1 in PBMCs Is Associated with Intracranial Aneurysm Formation, but TGFB3 Expression Implicated Rupture](#)
- [Risk Factors for Unfavorable Angiographic Outcomes after Reconstructive Endovascular Treatments of Unruptured Vertebral Artery Dissecting Aneurysms](#)
- [The role of systemic inflammation in the formation and rupture of intracranial aneurysms in moyamoya disease: a retrospective cohort study](#)
- [Risk factors for rupture of intracranial aneurysms in patients with autoimmune diseases](#)
- [Association between triglyceride-glucose index and intracranial aneurysm rupture: findings from a retrospective study](#)
- [Development and validation of a nomogram for predicting intracranial infection after intracranial aneurysm surgery](#)
- [Apolipoprotein A1 Levels Are Inversely Associated with Aneurysm Wall Enhancement in Unruptured Intracranial Aneurysms](#)

see [Aneurysm risk factors](#).

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=intracranial_aneurysm_risk_factors

Last update: **2024/06/07 02:50**

