

Intrajejunal levodopa-carbidopa therapy

Intrajejunal [levodopa-carbidopa](#) therapy and an [apomorphine](#) pump can be considered for advanced PD with fluctuations not sufficiently managed with oral treatments ^{1) 2)}.

1)

Deuschl G, Antonini A, Costa J, Śmiłowska K, Berg D, Corvol JC, Fabbrini G, Ferreira J, Foltynie T, Mir P, Schrag A, Seppi K, Taba P, Ruzicka E, Selikhova M, Henschke N, Villanueva G, Moro E. European Academy of Neurology/Movement Disorder Society-European Section Guideline on the Treatment of Parkinson's Disease: I. Invasive Therapies. *Mov Disord*. 2022 Jul;37(7):1360-1374. doi: 10.1002/mds.29066. Epub 2022 Jul 6. PMID: 35791767.

2)

Deuschl G, Antonini A, Costa J, Śmiłowska K, Berg D, Corvol JC, Fabbrini G, Ferreira J, Foltynie T, Mir P, Schrag A, Seppi K, Taba P, Ruzicka E, Selikhova M, Henschke N, Villanueva G, Moro E. European Academy of Neurology/Movement Disorder Society - European Section guideline on the treatment of Parkinson's disease: I. Invasive therapies. *Eur J Neurol*. 2022 Sep;29(9):2580-2595. doi: 10.1111/ene.15386. Epub 2022 Jul 6. PMID: 35791766.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=intrajejunal_levodopa-carbidopa_therapy

Last update: **2024/06/07 02:50**

