

Lactic acidosis

Lactic [acidosis](#) is a form of [metabolic acidosis](#) that begins when a person overproduces or underutilizes [lactic acid](#), and their body is not able to adjust to these changes.

Generally, lactate levels > 2 mmol/L represent hyperlactatemia, whereas [lactic acidosis](#) is often defined as lactate > 4 mmol/L.

Although hyperlactatemia is common finding in [liver transplant](#) (LT) candidates, association between lactate and organ failures with Acute-on-chronic Liver Failure (ACLF) is poorly studied.

A total of 2,002 patients from LT registry between January 2008 and February 2019 were analyzed. Six organ failures (liver, kidney, brain, coagulation, circulation, and lung) were defined by criteria of EASL-CLIF ACLF Consortium. Variable importance of preoperative hyperlactatemia was examined by machine learning using random survival forest (RSF). Kaplan-Meier Survival curve analysis was performed to assess 90-day mortality.

Median lactate level was 1.9 mmol/L (interquartile range: 1.4, 2.4 mmol/L) and 107 (5.3%) patients showed > 4.0 mmol/L. RSF analysis revealed that the four most important variables for hyperlactatemia were MELD score, circulatory failure, hemoglobin, and respiratory failure. The 30-day and 90-day mortality rates were 2.7% and 5.1%, whereas patients with lactate > 4.0 mmol/L showed increased rate of 15.0% and 19.6%, respectively.

About 50% and 5% of LT candidates showed pre-LT hyperlactatemia of > 2.0 mmol/L and > 4.0 mmol/L, respectively. Pre-LT lactate > 4.0 mmol/L was associated with increased early post-LT mortality. Our results suggest that future study of correcting modifiable risk factors may play a role in preventing hyperlactatemia and lowering early mortality after LT ¹⁾

Risk factors

[Iodinated contrast](#) (IV or intra-arterial) may delay the excretion of [metformin](#) (Glucophage®, Avandamet®), an oral hypoglycemic agent used in [type 2 diabetes](#), and can be associated with [lactic acidosis](#) and [renal failure](#) (particularly in patients with CHF or those consuming [alcohol](#)). The manufacturer recommends withholding metformin 48 hrs prior to and following [contrast](#) administration (or longer if there is evidence of declining renal function following the use of contrast). Metformin should also be held ≈ 48 hours before any surgery, and should not be restarted post-op until the patient has fully recovered and is eating and drinking normally. The maximum dose of iodine with normal renal function is ≈ 86 gm in a 24-hour period.

Complications

People with lactic acidosis have problems with their [liver](#) (and sometimes their kidneys) being able to remove excess acid from their body.

Coma.

Cheong et al. searched the important variables for pre-LT hyperlactatemia and examined the impact of preoperative hyperlactatemia on early mortality after LT.

A total of 2,002 patients from LT registry between January 2008 and February 2019 were analyzed. Six organ failures (liver, kidney, brain, coagulation, circulation, and lung) were defined by criteria of EASL-CLIF ACLF Consortium. Variable importance of preoperative hyperlactatemia was examined by machine learning using random survival forest (RSF). Kaplan-Meier Survival curve analysis was performed to assess 90-day mortality.

Median lactate level was 1.9 mmol/L (interquartile range: 1.4, 2.4 mmol/L) and 107 (5.3%) patients showed > 4.0 mmol/L. RSF analysis revealed that the four most important variables for hyperlactatemia were MELD score, circulatory failure, hemoglobin, and respiratory failure. The 30-day and 90-day mortality rates were 2.7% and 5.1%, whereas patients with lactate > 4.0 mmol/L showed increased rate of 15.0% and 19.6%, respectively.

About 50% and 5% of LT candidates showed pre-LT hyperlactatemia of > 2.0 mmol/L and > 4.0 mmol/L, respectively. Pre-LT lactate > 4.0 mmol/L was associated with increased early post-LT mortality. Results suggest that future study of correcting modifiable risk factors may play a role in preventing hyperlactatemia and lowering early mortality after LT ²⁾.

1) , 2)

Cheong Y, Lee S, Lee DK, Kim KS, Sang BH, Hwang GS. Preoperative hyperlactatemia and early mortality after liver transplantation: selection of important variables using random forest survival analysis. *Anesth Pain Med* (Seoul). 2021 Oct;16(4):353-359. doi: 10.17085/apm.21049. Epub 2021 Oct 14. PMID: 35139616.

From:
<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:
https://neurosurgerywiki.com/wiki/doku.php?id=lactic_acidosis

Last update: **2024/06/07 02:51**

