

En el siglo XIX, John Hunter demostró que la ligadura proximal de arterias periféricas era un método confiable para tratar aneurismas periféricos. Esto originó la “ligadura hunteriana”.

Siglos posteriores, [Víctor Horsley](#) y [Harvey Cushing](#) extendieron esta técnica a los [aneurismas cerebrales](#), con un menor éxito debido a la tasa de mortalidad de hasta un 25 por ciento.

Aunque el uso de la oclusión de la arteria proximal, o ligadura Hunteriana, para el tratamiento de aneurismas intracraneales ha disminuido mucho durante las últimas décadas, este enfoque todavía encuentra uso para ciertos [aneurismas gigantes](#) y [complejos](#). Los principales riesgos de sacrificio de esta arteria son complicaciones [isquémicas](#) pero también, aunque rara, formación de aneurismas de novo (Arnaout y col., 2012).

Bibliografía

Arnaout, Omar M, Rudy J Rahme, Salah G Aoun, Marc R Daou, H Hunt Batjer, and Bernard R Bendok. 2012. “De Novo Large Fusiform Posterior Circulation Intracranial Aneurysm Presenting with Subarachnoid Hemorrhage 7 Years After Therapeutic Internal Carotid Artery Occlusion: Case Report and Review of the Literature.” *Neurosurgery* 71 (3) (September): E764–771.
doi:10.1227/NEU.0b013e31825fd169.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=ligadura_hunteriana

Last update: **2025/03/10 15:15**

