

Lumbar puncture for idiopathic normal pressure hydrocephalus diagnosis

- [Diagnostic efficacy of radionuclide scintigraphy in detecting lumboperitoneal shunt obstructions in idiopathic hydrocephalus and intracranial hypertension](#)
- [Neurocognitive effects of CSF biomarkers in idiopathic normal pressure hydrocephalus patients undergoing VP shunt placement](#)
- [Determinants of clinical response in possible normal pressure hydrocephalus](#)
- [Progressive ataxia, cognitive decline, urinary incontinence, and unexplained hydrocephalus: a rare case of idiopathic normal pressure hydrocephalus in epileptic patient](#)
- [Long-term gait improvement following a CSF tap test in idiopathic normal pressure hydrocephalus patients: an analysis of clinical outcomes](#)
- [Temporary Drainage of Cerebrospinal Fluid for Diagnosis and Treatment of Hydrocephalus](#)
- [Standardizing the large-volume "tap test" for evaluating idiopathic normal pressure hydrocephalus: a systematic review](#)
- [Treatment of INPH: novel insights](#)

Opening pressure

see [Opening pressure](#).

Lumbar infusion test

see [Lumbar infusion test for idiopathic normal pressure hydrocephalus diagnosis](#).

Cerebrospinal fluid tap test

[Cerebrospinal fluid tap test](#) (CSF-TT), are often used in practice to provide further predictive value in detecting suitable patients for shunting.

Intraventricular infusion test

[Intraventricular infusion test](#)

From:
<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:
https://neurosurgerywiki.com/wiki/doku.php?id=lumbar_puncture_for_idiopathic_normal_pressure_hydrocephalus_diagnosis

Last update: **2024/06/07 02:59**

