2025/05/10 18:26 1/1 2016

see Intracranial meningioma case series

see Spinal meningioma case series

2016

Gousias et al., reviewed their institutional experience with a policy based on maximal safe resections for meningiomas, and they analyzed the impact of the degree of resection on functional outcome and progression free survival (PFS).

They retrospectively analyzed 901 consecutive patients with primary meningiomas (716 WHO Grade I, 174 Grade II, and 11 Grade III) who underwent resections at the University Hospital of Bonn between 1996 and 2008. Clinical and treatment parameters as well as tumor characteristics were analyzed using standard statistical methods.

The median follow-up was 62 months. PFS rates at 5 and 10 years were 92.6% and 86.0%, respectively. Younger age, higher preoperative Karnofsky Performance Scale (KPS) score, and convexity tumor location, but not the degree of resection, were identified as independent predictors of a good functional outcome (defined as KPS Score 90-100). Independent predictors of PFS were degree of resection (Simpson Grade I vs II vs III vs IV), MIB-1 index (< 5% vs 5%-10% vs > 10%), histological grade (WHO I vs II vs III), tumor size (≤ 6 vs > 6 cm), tumor multiplicity, and location. A Simpson Grade II rather than Grade I resection more than doubled the risk of recurrence at 10 years in the overall series (18.8% vs 8.5%). The impact of aggressive resections was much stronger in higher grade meningiomas.

A policy of maximal safe resections for meningiomas prolongs PFS and is not associated with increased morbidity ¹⁾.

1)

Gousias K, Schramm J, Simon M. The Simpson grading revisited: aggressive surgery and its place in modern meningioma management. J Neurosurg. 2016 Sep;125(3):551-60. doi: 10.3171/2015.9.JNS15754. Epub 2016 Jan 29. PubMed PMID: 26824369.

From:

https://neurosurgerywiki.com/wiki/ - Neurosurgery Wiki

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=meningioma_case_series

Last update: 2025/04/29 20:29

