

Microcystic meningioma diagnosis

- [Neuroradiological Evaluation of Anatomic-Morphometric Arcuate Fascicle Modifications According to Different Brain Tumor Histotypes: An Italian Multicentric Study](#)
 - [Mixed reality holographic navigation for intracranial lesions using HoloLens 2: A pilot study and literature review](#)
 - [Advancing Neurosurgical Oncology and AI Innovations in Latin American Brain Cancer Care: Insights from a Center of Excellence](#)
 - [Investigation of the frequency of meningotheelial hyperplasia and its clinicopathological correlation in patients diagnosed with subdural hematoma](#)
 - [Cerebral Lymphangioma: A Clinical Case History](#)
 - [Predicting intraoperative meningioma consistency using features from standard MRI sequences: a preoperative evaluation](#)
 - [The dural tail in intracranial meningioma: Heads up or tail down? A systematic review of the literature](#)
 - [Simultaneous surgical management of a giant tuberculum sellae meningioma and pregnancy-related complications: a case report and literature review](#)
-
-

1. Clinical evaluation

Symptoms depend on tumor location: headaches, seizures, focal deficits, or signs of increased intracranial pressure.

Sometimes found incidentally on imaging.

2. Imaging

MRI Characteristics

Extra-axial, well-circumscribed mass.

T2-weighted images: very high signal intensity (bright), more than typical meningiomas, due to the high fluid content of microcysts.

T1-weighted images: usually iso- to hypointense.

After contrast (gadolinium), enhancement can be heterogeneous (not uniform) — different from classic meningiomas which often enhance homogeneously.

May show a “soap-bubble” or reticular appearance.

Sometimes associated with a dural tail.

SPECT

Microcystic [meningiomas](#) are rare but benign brain tumors. Previous reports have shown that Thallium-201 single-photon emission computed tomography (201Tl SPECT) demonstrated a higher late-phase accumulation of 201Tl in malignant or recurrent meningiomas than in nonaggressive meningiomas.

A case of a microcystic meningioma with a high 201Tl SPECT retention rate in a 62-year-old woman who complained of headache. Computed tomography revealed an intracranial tumor in the right frontal lobe. Moreover, 201Tl SPECT revealed a high uptake of 201Tl in the tumor, which was particularly prominent in the delayed phase. The uptake index on an early image was 1.46 and that on a delayed image was 1.35. Therefore, the retention index was 0.92. After 2 years of tumor growth, we performed successful radical resection, and histological examination revealed the presence of a microcystic meningioma. Therefore, 201Tl SPECT may be useful for the preoperative diagnosis of microcystic meningiomas and that late-phase accumulation of 201Tl is not a specific finding of malignant brain tumors. Therefore, we need to be careful in the evaluation and judgment of high retention in a delayed image of 201Tl SPECT ¹⁾.

Fluciclovine Positron Emission Tomography

A 69-year-old asymptomatic woman underwent [magnetic resonance imaging](#) (MRI), revealing a left [parietal lobe tumor](#) suspected to be a [high-grade glioma](#) (HGG). [18F][Fluciclovine positron emission tomography](#) (PET), which reflects amino acid metabolism, showed moderate uptake [SUVmax 1.4, tumor-to-normal ratio (TNR) 1.9], consistent with HGG, whereas [18F][fluorodeoxyglucose](#)-PET findings demonstrated low uptake (SUVmax 2.5, TNR 0.22), which is atypical for HGG. Surgical and pathologic investigations confirmed the diagnosis of [microcystic meningioma](#), a rare subtype classified as grade 1. [Microcystic meningiomas](#) mimic HGG on magnetic resonance imaging and [18F]fluciclovine-PET due to L-amino acid transporter expression, complicating diagnosis ²⁾

This report compellingly illustrates that fluciclovine uptake is not specific for high-grade malignancy in CNS tumors. Clinicians must interpret [PET](#) findings in conjunction with [clinical history](#), MRI features, and alternative PET tracers like FDG. Future work should include larger studies evaluating false-positive rates of fluciclovine-PET in [brain tumors](#) and the molecular basis of tracer uptake in benign lesions like meningiomas.

3. Histopathology

Essential for definitive diagnosis.

Microscopic features:

Loose, edematous tissue.

Abundant microcysts filled with fluid.

Round/oval nuclei, sparse mitotic activity.

Immunohistochemistry

Positive for EMA (epithelial membrane antigen).

Positive for vimentin.

Low Ki-67 (proliferation index), consistent with low-grade tumor.

4. Differential diagnosis

Other cystic brain tumors (e.g., hemangioblastoma, pilocytic astrocytoma, metastases).

Imaging and pathology together help distinguish it.

1)

Matano F, Adachi K, Murai Y, Kitamura T, Ohashi R, Teramoto A, Morita A. Microcystic Meningioma with Late-phase Accumulation on Thallium-201 single-photon emission computed tomography: Case Report. *Neurol Med Chir (Tokyo)*. 2014 Jan 10. [Epub ahead of print] PubMed PMID: 24418788.

2)

Takenaka J, Hirata K, Watanabe S, Ishi Y, Kudo K. Fluciclovine-PET Uptake in Microcystic Meningioma Mimicking High-grade Glioma: A Case Report. *Clin Nucl Med*. 2025 Apr 24. doi: 10.1097/RLU.0000000000005924. Epub ahead of print. PMID: 40279665.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=microcystic_meningioma_diagnosis

Last update: **2025/04/26 09:54**

