Middle cerebral artery aneurysm etiology

- Effectiveness of Decompressive Craniectomy for Severe Cerebral Infarction Caused by Stanford Type A Acute Aortic Dissection:Report of a Case
- Infected aneurysm of middle meningeal artery causing intracerebral hemorrhage: illustrative case
- The role of systemic inflammation in the formation and rupture of intracranial aneurysms in moyamoya disease: a retrospective cohort study
- Protective Effect of Resveratrol Against Intracranial Aneurysm Rupture in Mice
- Association between triglyceride-glucose index and intracranial aneurysm rupture: findings from a retrospective study
- Outcomes of flow diversion in intracranial aneurysms accompanied by parent artery stenosis
- External Validation of the ARISE Prediction Models for Aneurysmal Rebleeding After Aneurysmal Subarachnoid Hemorrhage
- When an aneurysm pretends to be a tumor: thrombosed posterior inferior cerebellar artery aneurysm mimicking a neoplasm in a pediatric patient. Illustrative case

Middle cerebral artery aneurysm formation is strongly associated with blood vessel tortuosity parameters, which can potentially be used to screen for patients at risk for MCA aneurysm formation [1].

[1] Kliś KM, Krzyżewski RM, Kwinta BM, Stachura K, Moskała M, Tomaszewski KA. Computer-aided analysis of middle cerebral artery tortuosity: association with aneurysm development. J Neurosurg. 2018 May 18:1-7. doi: 10.3171/2017.12.JNS172114. [Epub ahead of print] PubMed PMID: 29775150.

From:
https://neurosurgerywiki.com/wiki/ - **Neurosurgery Wiki**

Permanent link:
https://neurosurgerywiki.com/wiki/doku.php?id=middle_cerebral_artery_aneurysm_etiology

Last update: **2024/06/07 02:52**