

Neurosyphilis treatment

- [High-Dose Benzylpenicillin Treatment-Induced Febrile Neutropenia in HIV-Infected Male with Neurosyphilis: Case Report](#)
- [Syphilitic Aortitis with Concomitant Neurosyphilis in Asymptomatic Patient](#)
- [Secondary syphilis presenting with increased ophthalmic artery resistance index: A case report](#)
- [Optic Neuritis As the Initial Presentation of Syphilis](#)
- [Hypertensive Disc Edema or Ocular Syphilis? A Case Report of the Great Masquerader](#)
- [Patients with late syphilis and neurosyphilis treated in Bialystok in 2014-2023](#)
- [Comparative analysis of language models in addressing syphilis-related queries](#)
- [Unraveling an Enigmatic Triad: A Case Report of Concurrent Neurosyphilis, Ocular Syphilis, and Ootosyphilis in a Patient with HIV](#)

Neurosyphilis treatment usually consists of intravenous antibiotics, such as [penicillin](#), administered over a course of time. In some cases, additional medications or interventions may be necessary to manage symptoms or complications.

Given the serious nature of neurosyphilis and its potential long-term consequences, it's crucial for individuals with syphilis to undergo regular testing and to seek treatment promptly if diagnosed. Preventive measures, such as practicing safe sex and using condoms, can also help reduce the risk of contracting syphilis and its complications.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=neurosyphilis_treatment

Last update: **2024/06/07 02:55**

