


Olfactory groove meningioma magnetic resonance imaging



Extraaxial space-occupying lesion measuring 3.8 x 3.8 x 2.6 cm with a dural base on the [planum sphenoidale](#) and [olfactory groove](#) compatible with [planum sphenoidale meningioma](#) or [olfactory groove meningioma](#). Most of the lesion shows intense [enhancement](#), with some areas without enhancement attributable to [cystic-necrotic](#) changes. [Diffusion](#) shows a peripheral component with low values on the [ADC](#) map and a central component with high values that suggest [hypercellularity](#) and hypocellularity, respectively. [Dural tail](#) extending to the [tuberculum sellae](#) is also seen.

It conditions extensive [edema](#) in the [white matter](#) of both [frontal lobes](#) and knee of the [corpus callosum](#) and obliteration of the frontal grooves.

Associated [sclerosis](#) of the [sphenoid sinus](#). No bone or brain [invasion](#) is seen. The posterior margin of the lesion is adjacent to the intracranial segment of both [optic nerves](#), with no compression of the [optic pathway](#). The A2 segment of both [anterior cerebral artery](#) contacts the posterior margin of the lesion. [Ventricular system](#): normal size and morphology for the patient's age. Midline and basal cisternae: normal.

Chronic ischemic lesions in the lower region of the left cerebellar hemisphere (territory of the left posterior inferior cerebellar artery). Preserved arterial flow vacuum.

From:
<https://neurosurgerywiki.com/wiki/> - Neurosurgery Wiki

Permanent link:
https://neurosurgerywiki.com/wiki/doku.php?id=olfactory_groove_meningioma_magnetic_resonance_imaging

Last update: **2024/06/07 02:57**

