

Perimesencephalic Subarachnoid Hemorrhage Epidemiology

- [Subarachnoid hemorrhage part 1 : Pathophysiology, diagnostics and special forms](#)
- [Association of SSRI and SNRI use with incident hyponatremia after subarachnoid hemorrhage](#)
- [Risk stratification of delayed causative aneurysm detection and long-term outcome in angiographically negative spontaneous subarachnoid haemorrhage](#)
- [Cognitive, Mental Health, Functional, and Quality of Life Outcomes 1 Year After Spontaneous Subarachnoid Hemorrhage: A Prospective Observational Study](#)
- [Treatment and outcomes of non-aneurysmal perimesencephalic subarachnoid haemorrhage: A 5 year retrospective study in a tertiary care centre](#)
- [Physical exertion as a risk factor for perimesencephalic nonaneurysmal subarachnoid hemorrhage](#)
- [Cannabis Use and Increased Nonaneurysmal Subarachnoid Hemorrhage in the Past Decade](#)
- [Clinical comparison of patients with angiography: Negative perimesencephalic and nonperimesencephalic subarachnoid hemorrhages](#)

[Perimesencephalic subarachnoid hemorrhage](#) is rare with an [incidence](#) of 0.5 in 100 000 in adults.

PMSAH represents 5-10% of all [subarachnoid haemorrhages](#) and ~33% of all non-aneurysmal SAH.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=perimesencephalic_subarachnoid_hemorrhage_epidemiology

Last update: **2024/06/07 02:55**

