

Posterior inferior cerebellar artery aneurysm surgery

In 1947, Rizzoli and Hayes trapped and excised a [Posterior inferior cerebellar artery aneurysm](#)¹⁾.

In recent years, the trend of treatment of [posterior circulation aneurysms](#) has tilted toward the endovascular arm.

[Posterior inferior cerebellar artery aneurysm](#) is challenging due to the deep location and intimate relation with the [medulla](#) and [cranial nerves](#): [glossopharyngeal nerves](#) (IX), [vagus nerves](#) (X), [accessory nerves](#) (XI).

Microsurgery remains the bedrock for treating PICA aneurysms, with planning depending on size, location, shape, presentation, and clinical condition of the patient.

Approaches

PICA has the most complex and variable course among all arteries of the posterior circulation. The complexities involved in the surgical approach to the PICA aneurysms relate to the narrow corridor limited by the [brainstem](#), petrous-occipital bones, and multiple neurovascular structures occupying the [cerebellomedullary cistern](#) and [cerebellopontine cisterns](#)²⁾.

Surgical approach is influenced by VA-BA tortuosity and variations in anatomy, location of the VA-BA junction and the PICA aneurysm relative to the brainstem, and the pattern of collateral supply. The special category of VA-PICA junctional aneurysms and its management; and, the multiple anatomical variations of PICA aneurysms, merit special surgical considerations and have been highlighted in the study of Singh et al.³⁾

The [far lateral transcondylar craniotomy](#) is the standard approach for [posterior inferior cerebellar artery aneurysm](#) exposure through the microsurgical dissection in the [vagoaccessory triangle](#) (VAT). However, the extended [retrosigmoid craniotomy](#) and dissection through the [glossopharyngeal-cochlear triangle](#) (GCT) may be more appropriate when the patient has an aneurysm arising from a high-riding [vertebral artery](#) (VA)-PICA origin.

Midline approaches are safe, familiar, and more reasonable, especially for distal PICA aneurysms. Neurovascular conflicts, proximal perforators, and [eloquent areas](#) influence dissection and [clipping](#) techniques. Options include direct [clipping](#), clipping with [wrapping](#), wrapping, resection, proximal occlusion or trapping with revascularization, and distal occlusion⁴⁾.

In aneurysms that cannot be clipped, sacrifice of the PICA without revascularisation procedures in proximally located PICA aneurysms may still be feasible if the occlusion is done distal to the

perforators ⁵⁾.

Far lateral approach

Although for anterior medullary segment, [far lateral approach](#) provides the best exposure to the [cerebellomedullary fissure](#), the recent trend has been on a tailored approach. Seoane et al ⁶⁾ reported a surgical series using the far-lateral approach to PICA aneurysms without [condylar resection](#). Rodríguez-Hernández and Lawton described the vagoaccessory triangle that is bordered medially by the medulla, laterally by the spinal accessory nerve (CN XI), and superiorly by the vagus nerve (CN X), as an anatomical corridor to reach the PICA aneurysms. ⁷⁾ As for distal PICA aneurysms, midline approach is, without doubt, sufficient. In the series, of Deora et al. ⁸⁾ they used the [far lateral approach](#) with condylar resection only in 25% (5/20) of the cases, with the majority (4/5) being anterior medullary or VA-PICA junction cases with acceptable complication rates. Other improvised techniques have been described such as the [lateral suboccipital approach](#), the lateral suboccipital [transcondylar approach](#), removal of the posterior condyle, C1 lateral mass, and/or the jugular tubercle, all of which help in widening the surgical corridor. ⁹⁾ Proximal control of VA can be secured by the far-lateral approach and dissection of the arachnoid caudal to the IX and X CNs, avoiding the resection of the posterior lip of the [foramen magnum \(opisthion\)](#) ¹⁰⁾.

Other more complex skull-base approaches have also been described, such as [presigmoid retrolabyrinthine approach](#) and [transsigmoid approach](#). A modified far lateral approach, which has been the approach most frequently used by Ng Hua Bak Ivan in dealing with PICA aneurysms, is described in https://link.springer.com/chapter/10.1007/978-981-10-8950-3_19#Sec3

Videos

The [far lateral transcondylar craniotomy](#) is the standard approach for [posterior inferior cerebellar artery aneurysm](#) exposure through the microsurgical dissection in the [vagoaccessory triangle](#) (VAT). However, the [extended retrosigmoid craniotomy](#) and dissection through the [glossopharyngeal-cochlear triangle](#) (GCT) may be more appropriate when the patient has an aneurysm arising from a high-riding [vertebral artery](#) (VA)-PICA origin.

Peitz et al. presented a case of a 41-yr-old woman with hypertension presenting with left occipital pain and left-side hearing loss and past facial spasm and pain. Computed tomography angiography and digital subtraction angiography demonstrated an unruptured 8.4 × 9.0 × 10.2 mm saccular aneurysm at the left VA-PICA junction. Surgical clipping was chosen over endovascular therapy given the relationship of the PICA origin to the aneurysm neck as well as the history of cranial [neuropathy](#). It was noted that the VA-PICA junction and aneurysm were high-riding at the level of the [internal auditory canal](#). An eRS craniotomy was performed with dissection through the GCT, and the aneurysm was clipped as shown in the accompanying 2-dimensional operative video. Postoperative angiography demonstrated complete occlusion of the aneurysm and patency of the left VA and PICA without stenosis, and the patient had a favorable postoperative course although her left-sided hearing remained diminished. The eRS craniotomy allowed direct exposure via the GCT for clipping of the high-riding VA-PICA junction aneurysm and decompression of the cranial nerves. The traditional FL

craniotomy and exposure through the VAT would likely have resulted in a less desirable inferior trajectory.¹¹⁾

Occipital artery to posterior inferior cerebellar artery bypass for posterior inferior cerebellar artery aneurysm treatment

[Occipital artery to posterior inferior cerebellar artery bypass for posterior inferior cerebellar artery aneurysm treatment](#)

References

1)

Rizzoli H V, Hayes G J. Congenital berry aneurysm of the posterior fossa; case report with successful operative excision. J Neurosurg. 1953;10(05):550-551.

2)

Rodríguez-Hernández A, Lawton MT. Anatomical triangles defining surgical routes to posterior inferior cerebellar artery aneurysms. J Neurosurg. 2011;114:1088-94.

3)

Singh RK, Behari S, Kumar V, Jaiswal AK, Jain VK. Posterior inferior cerebellar artery aneurysms: Anatomical variations and surgical strategies. Asian J Neurosurg. 2012 Jan;7(1):2-11. doi: 10.4103/1793-5482.95687. PMID: 22639684; PMCID: PMC3358952.

4)

Lehto H, Harati A, Niemelä M et al. Distal posterior inferior cerebellar artery aneurysms: clinical features and outcome of 80 patients. World Neurosurg. 2014;82(05):702-713.

5)

Liew D, Ng PY, Ng I. Surgical management of ruptured and unruptured symptomatic posterior inferior cerebellar artery aneurysms. Br J Neurosurg. 2004 Dec;18(6):608-12. doi: 10.1080/02688690400022870. PMID: 15799193.

6)

Seoane P, Kalb S, Clark J C et al. Far-lateral approach without drilling the [occipital condyle](#) for vertebral artery-posterior inferior cerebellar artery aneurysms. Neurosurgery. 2017;81(02):268-274.

7)

Rodríguez-Hernández A, Lawton M T. Anatomical triangles defining surgical routes to posterior inferior cerebellar artery aneurysms. J Neurosurg. 2011;114(04):1088-1094.

8) 10)

Deora H, Nayak N, Dixit P, Vikas V, Rao KVLN, Pruthi N, Srinivas D, Shukla DP, Bhat DI, Malla BR, Devi BI, Somanna S. Surgical Management and Outcomes of Aneurysms of Posterior Inferior Cerebellar Artery: Location-Based Approaches with Review of Literature. J Neurosci Rural Pract. 2020 Jan;11(1):34-43. doi: 10.1055/s-0039-3399486. Epub 2020 Jan 27. PMID: 32269450; PMCID: PMC7138643.

9)

Viswanathan G C, Menon G, Nair S, Abraham M. Posterior inferior cerebellar artery aneurysms: operative strategies based on a surgical series of 27 patients. Turk Neurosurg. 2014;24(01):30-37.

11)

Peitz GW, McDermott RA, Baranoski JF, Lawton MT, Mascitelli JR. Extended Retrosigmoid Craniotomy and Approach Through the Glossopharyngeal Cochlear Triangle for Clipping of a High-Riding Vertebral-Posterior Inferior Cerebellar Artery Aneurysm: 2-Dimensional Operative Video. Oper Neurosurg (Hagerstown). 2021 May 14:opab140. doi: 10.1093/ons/opab140. Epub ahead of print.

PMID: 33989426.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=posterior_inferior_cerebellar_artery_aneurysm_surgery

Last update: **2024/06/07 02:53**

