

RNA synthesis

Synthesis of **RNA** is usually catalyzed by an enzyme—**RNA polymerase**—using **DNA** as a template, a process known as **transcription**. Initiation of transcription begins with the binding of the enzyme to a promoter sequence in the DNA (usually found “upstream” of a gene). The DNA double helix is unwound by the **helicase** activity of the enzyme. The enzyme then progresses along the template strand in the 3′ to 5′ direction, synthesizing a **complementary RNA** molecule with elongation occurring in the 5′ to 3′ direction. The DNA sequence also dictates where termination of RNA synthesis will occur.

Primary transcript RNAs are often modified by enzymes after transcription. For example, a poly(A) tail and a 5' cap are added to eukaryotic pre-mRNA and introns are removed by the spliceosome.

There are also a number of RNA-dependent RNA polymerases that use RNA as their template for synthesis of a new strand of RNA. For instance, a number of RNA viruses (such as poliovirus) use this type of enzyme to replicate their genetic material.

Also, RNA-dependent RNA polymerase is part of the **RNA interference** pathway in many organisms.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=rna_synthesis

Last update: **2024/06/07 02:50**

