

Thalamic Diffuse astrocytoma

- [Minimally Invasive and Cost-Effective Access to Deep-Seated Intracranial Lesions Using 19F Peel-Away Sheath Introducer and "Dynamic" Retraction: Technical Note and a Case Series](#)
- [Stereotactic laser ablation for pediatric central nervous system tumors: a systematic review and meta-analysis of the literature](#)
- [High-grade astrocytoma with piloid features: a single-institution case series and literature review](#)
- [Characterizing the molecular and spatial heterogeneity of midline gliomas in adults: a single institution analysis](#)
- [High-grade astrocytoma in a sheep: clinical, pathology, and genetic investigations](#)
- [BRAF inhibitor monotherapy in BRAFV600E-mutated pediatric low-grade glioma: a single center's experience](#)
- [Free-hand Frameless Pinless Electromagnetic-Navigation \(AXIEM™\)-Guided Brain Lesion Biopsies: An Institution Based Experience from a Low-Middle-Income Country](#)
- [A Case of Anterior Cingulotomy for Intractable Pain Caused Thalamic Glioma](#)

Thalamic [Diffuse astrocytoma](#), IDH-mutant (WHO Grade 2) are rare gliomas that arise in the thalamus,

These tumors belong to the category of diffuse gliomas, which infiltrate surrounding brain tissue and typically exhibit slow growth. However, they carry a significant clinical burden due to their deep location and the challenges in surgical resection.

Epidemiology

Most commonly seen in children and young adults.

Can also occur in adults but are less frequent compared to cerebral hemispheric gliomas.

No significant sex predilection.

From:
<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:
https://neurosurgerywiki.com/wiki/doku.php?id=thalamic_diffuse_astrocytoma

Last update: **2025/03/03 10:41**

