

# Traumatic intracranial pseudoaneurysm

Traumatic intracranial pseudoaneurysms are a rare but severe complication following arterial injury. Pseudoaneurysm formation can occur secondary to blunt or penetrating trauma or iatrogenic injury. We report a case of traumatic pseudoaneurysm secondary [intracranial pressure monitoring](#). A 27-year-old man was involved in a motorcycle accident resulting in multiple intracranial hemorrhages. The patient underwent craniectomy and placement of an ICP monitor. 17 days later he developed dilation of his left pupil, with imaging demonstrating a new hemorrhage in the vicinity of the previous ICP monitor. A cerebral angiogram confirmed a left-sided distal M4 pseudoaneurysm which was treated by n-butyl cyanoacrylate embolization. Intracranial pseudoaneurysm formation following neurosurgical procedures is uncommon. Delayed intracranial hemorrhage in a region of prior intracranial manipulation, even following a procedure as 'routine' as placement of an ICP monitor, should raise the suspicion for this rare but potentially lethal complication <sup>1)</sup>

<sup>1)</sup>

Shah KJ, Jones AM, Arnold PM, Ebersole K. Intracranial pseudoaneurysm after intracranial pressure monitor placement. J Neurointerv Surg. 2014 Dec 17. pii: neurintsurg-2012-011410.rep. doi: 10.1136/neurintsurg-2014-011410.rep. [Epub ahead of print] PubMed PMID: 25520265.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

[https://neurosurgerywiki.com/wiki/doku.php?id=traumatic\\_intracranial\\_pseudoaneurysm](https://neurosurgerywiki.com/wiki/doku.php?id=traumatic_intracranial_pseudoaneurysm)

Last update: **2025/03/10 15:19**

