

Treatment refusal

- Middle meningeal artery (MMA) embolisation for chronic subdural haematomas: rationale and design for the STOp Recurrence of MMA Bleeding (STORMM) randomised control trial-a study protocol
- The Safety and Feasibility of a Care Hotel Model in Elective Neuroendovascular Interventions: A Pilot Study
- Long-term survival and undetectable circulating tumor DNA following comprehensive involved site radiotherapy for oligometastases
- Disparities in consent to adjuvant radiotherapy in primary glioblastoma: a population-based study
- Exploring the perspectives of early-stage postoperative glioblastoma patients and their caregivers on end-of-life care planning: An exploratory qualitative interview study
- Association between COVID-19 anxiety syndrome and COVID-19 vaccine hesitancy in the postpandemic era: a cross-sectional study in Hong Kong
- A 5-year Review of Characteristics and Outcomes of Trauma Surgery Patients Leaving Against Medical Advice
- Prognostic Factors in Idiopathic Sudden Sensorineural Hearing Loss: The Experience of Two Audiology Tertiary Referral Centres

In a study, patients with poor [Hunt and Hess Stroke Scale](#), [loss of consciousness](#) at symptom onset, or largest [intracranial aneurysm sizes](#) ≥ 5 mm in size showed a high [mortality rate](#). A high number of [treatment refusals](#) was present in this study. These findings have implications for medical insurance policy, doctor-patient communication, and popular science education ¹⁾

¹⁾

Yang Y, Wang B, Ge X, Lu W, Peng C, Zhao Y, Li J, Wang J, Lei P, Zhang J, Feng L, Liu Z, Yang X. Natural Course of Ruptured but Untreated Intracranial Aneurysms: A Multicenter 2-Year Follow-Up Study. Stroke. 2023 Jun 12. doi: 10.1161/STROKEAHA.123.042530. Epub ahead of print. PMID: 37306018.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**

Permanent link:

https://neurosurgerywiki.com/wiki/doku.php?id=treatment_refusal

Last update: **2024/06/07 02:57**

