

# Valsartan

Valsartan, sold under the brand name Diovan among others, is a medication used to treat [high blood pressure](#), heart failure, and diabetic kidney disease.

It belongs to a class of medications referred to as angiotensin II receptor blockers (ARBs). It is a reasonable initial treatment for high blood pressure.

It is taken by mouth.[8] Versions are available as the combination valsartan/hydrochlorothiazide, valsartan/amlodipine, valsartan/amlodipine/hydrochlorothiazide, or valsartan/sacubitril.


## Unclassified

- 1: Yang M, Zhang S, Liang J, Tang Y, Wang X, Huang C, Zhao Q. Different effects of norepinephrine and nerve growth factor on atrial fibrillation vulnerability. *J Cardiol*. 2019 Nov;74(5):460-465. doi: 10.1016/j.jcc.2019.04.009. Epub 2019 May 15. PubMed PMID: 31101572.
- 2: Sawano T, Ozaki A, Saito H, Shimada Y, Tanimoto T. Payments From Pharmaceutical Companies to Authors Involved in the Valsartan Scandal in Japan. *JAMA Netw Open*. 2019 May 3;2(5):e193817. doi: 10.1001/jamanetworkopen.2019.3817. Erratum in: *JAMA Netw Open*. 2019 Jul 3;2(7):e198619. PubMed PMID: 31099864; PubMed Central PMCID: PMC6537813.
- 3: Tsiofis C, Georgopoulos G, Oikonomou D, Thomopoulos C, Katsiki N, Kasiakogias A, Chrysochoou C, Konstantinidis D, Kalos T, Tousoulis D. Hypertension and Heart Failure with Preserved Ejection Fraction: Connecting the Dots. *Curr Vasc Pharmacol*. 2017;16(1):15-22. doi: 10.2174/1570161115666170414120532. Review. PubMed PMID: 28413968.
- 4: Nakamura K, Shimizu T, Yanagita T, Nemoto T, Taniuchi K, Shimizu S, Dimitriadis F, Yawata T, Higashi Y, Ueba T, Saito M. Angiotensin II acting on brain AT1 receptors induces adrenaline secretion and pressor responses in the rat. *Sci Rep*. 2014 Nov 28;4:7248. doi: 10.1038/srep07248. PubMed PMID: 25431019; PubMed Central PMCID: PMC5384216.
- 5: Wakai T, Yoshioka H, Yagi T, Kato T, Kinouchi H. Effects of valsartan on neuroprotection and neurogenesis after ischemia. *Neuroreport*. 2011 Jun 11;22(8):385-90. doi: 10.1097/WNR.0b013e328346be84. PubMed PMID: 21494160.
- 6: Aoki T, Nishimura M, Kataoka H, Ishibashi R, Miyake T, Takagi Y, Morishita R, Hashimoto N. Role of angiotensin II type 1 receptor in cerebral aneurysm formation in rats. *Int J Mol Med*. 2009 Sep;24(3):353-9. PubMed PMID: 19639227.
- 7: Nakazawa K, Ohta T, Fujimoto M, Imamura H, Hashimoto N. [Case report of cholesterol crystal embolism 1 month after carotid stenting]. *No Shinkei Geka*. 2007 Aug;35(8):807-11. Japanese. PubMed PMID: 17695780.
- 8: Madrid MI, García-Salom M, Tornel J, de Gasparo M, Fenoy FJ. Interactions between nitric oxide and angiotensin II on renal cortical and papillary blood flow. *Hypertension*. 1997 Nov;30(5):1175-82. PubMed PMID: 9369273.
- 9: Marano G, Argiolas L. Postjunctional regulation by angiotensin II of alpha 1-adrenoceptor-mediated

pressor responses in the rat. Eur J Pharmacol. 1994 Aug 11;261(1-2):121-6. PubMed PMID: 8001634.

From:

<https://neurosurgerywiki.com/wiki/> - **Neurosurgery Wiki**



Permanent link:

<https://neurosurgerywiki.com/wiki/doku.php?id=valsartan>

Last update: **2024/06/07 02:55**